

8. 消化器

(1) 上部消化管X線検査

① 1日人間ドック

<上部消化管X線検査判定結果>

<男性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	19	637	1,483	1,391	543	92	2	4,167	75.3
B1	2	26	140	168	56	15	0	407	7.4
B2	1	49	200	330	194	38	1	813	14.7
C	0	0	0	0	0	0	0	0	0.0
D	0	13	36	55	40	5	0	149	2.7
E	0	0	0	0	0	0	0	0	0.0
受診者数 (人)	22	725	1,859	1,944	833	150	3	5,536	100
有所見者数 (人)	3	88	376	553	290	58	1	1,369	24.7
有所見者率 (%)	13.6	12.1	20.2	28.4	34.8	38.7	33.3	24.7	

<女性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	6	377	860	750	307	47	0	2,347	67.2
B1	0	34	161	166	61	13	0	435	12.5
B2	1	62	174	278	119	15	0	649	18.6
C	0	0	0	0	0	0	0	0	0.0
D	0	6	16	26	6	5	0	59	1.7
E	0	0	0	0	0	0	0	0	0.0
受診者数 (人)	7	479	1,211	1,220	493	80	0	3,490	100.0
有所見者数 (人)	1	102	351	470	186	33	0	1,143	32.8
有所見者率 (%)	14.3	21.3	29.0	38.5	37.7	41.3	0.0	32.8	

<上部消化管X線検査疾患別一覧>

<男性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	総計
著変なし	19	637	1,483	1,391	543	92	2	4,167
慢性胃炎	1	29	122	223	149	32	1	557
びらん性胃炎	1	10	43	41	19	6	0	120
胃ポリープ	0	37	149	201	67	10	0	464
胃隆起性病変	0	4	9	15	10	1	0	39
胃潰瘍	0	0	0	1	1	0	0	2
胃潰瘍癒痕	0	4	10	23	17	3	0	57
胃粘膜下腫瘍	1	1	8	18	3	2	0	33
陰影斑(バリウム斑)	0	1	5	6	6	0	0	18
胃陥凹性病変	0	0	6	2	1	0	0	9
ひだ中断(粘膜中断)	0	1	1	0	1	0	0	3
粘膜不整	0	3	9	8	8	2	0	30
十二指腸潰瘍	0	0	0	0	0	0	0	0
十二指腸潰瘍癒痕	0	4	21	41	24	1	0	91
食道隆起性病変	0	1	5	14	5	0	0	25
総計	22	725	1,859	1,944	833	150	3	5,536

<女性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	総計
著変なし	6	377	860	750	307	47	0	2,347
慢性胃炎	1	19	69	130	73	10	0	302
びらん性胃炎	0	3	22	26	11	5	0	67
胃ポリープ	0	75	244	284	80	13	0	696
胃隆起性病変	0	2	6	10	5	4	0	27
胃潰瘍	0	0	0	1	0	0	0	1
胃潰瘍癒痕	0	0	2	4	2	0	0	8
胃粘膜下腫瘍	0	2	4	11	4	1	0	22
陰影斑(バリウム斑)	0	0	2	7	0	3	0	12
胃陥凹性病変	0	0	1	4	3	0	0	8

ひだ中断(粘膜中断)	0	1	0	0	0	0	0	1
粘膜不整	0	0	3	9	2	0	0	14
十二指腸潰瘍	0	1	0	0	0	0	0	1
十二指腸潰瘍癒痕	0	0	5	13	15	3	0	36
食道隆起性病変	0	1	3	0	2	0	0	6
総計	7	479	1,211	1,220	493	80	0	3,490

② 2日人間ドック

<上部消化管X線検査判定結果>

<男性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	0	3	14	10	7	2	0	36	73.5
B1	0	0	1	3	1	1	0	6	12.2
B2	0	1	2	2	1	1	0	7	14.3
C	0	0	0	0	0	0	0	0	0.0
D	0	0	0	0	0	0	0	0	0.0
E	0	0	0	0	0	0	0	0	0.0
受診者数 (人)	0	4	17	15	9	4	0	49	100.0
有所見者数 (人)	0	1	3	5	2	2	0	13	26.5
有所見者率 (%)	0.0	25.0	17.6	33.3	22.2	50.0	0.0	26.5	

<女性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	0	0	1	1	1	0	0	3	42.9
B1	0	0	0	1	1	0	0	2	28.6
B2	0	0	0	1	0	1	0	2	28.6
C	0	0	0	0	0	0	0	0	0.0
D	0	0	0	0	0	0	0	0	0.0
E	0	0	0	0	0	0	0	0	0.0
受診者数 (人)	0	0	1	3	2	1	0	7	100.0
有所見者数 (人)	0	0	0	2	1	1	0	4	57.1
有所見者率 (%)	0.0	0.0	0.0	66.7	50.0	100.0	0.0	57.1	

陰影斑(バリウム斑)	0	0	0	0	0	0	0	0
胃陥凹性病変	0	0	0	0	0	0	0	0
ひだ中断(粘膜中断)	0	0	0	0	0	0	0	0
粘膜不整	0	0	0	0	0	0	0	0
十二指腸潰瘍	0	0	0	0	0	0	0	0
十二指腸潰瘍癒痕	0	0	0	0	0	0	0	0
食道隆起性病変	0	0	0	0	0	0	0	0
総計	0	0	1	3	2	1	0	7

③ 施設内健診（協会けんぽ含む）

<上部消化管X線検査判定結果>

<男性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	17	1,222	1,922	1,122	668	53	0	5,004	65.1
B1	2	123	269	281	189	23	0	887	11.5
B2	3	189	441	404	389	19	0	1,445	18.8
C	0	0	0	0	0	0	0	0	0.0
D	0	26	89	119	107	14	0	355	4.6
E	0	0	0	0	0	0	0	0	0.0
受診者数 (人)	22	1,560	2,721	1,926	1,353	109	0	7,691	100.0
有所見者数 (人)	5	338	799	804	685	56	0	2,687	34.9
有所見者率 (%)	22.7	21.7	29.4	41.7	50.6	51.4	0.0	34.9	

<女性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	19	621	1,191	748	287	18	0	2,884	59.6
B1	1	117	369	273	118	13	0	891	18.4
B2	0	112	328	310	119	4	0	873	18.1
C	0	0	0	0	0	0	0	0	0.0
D	0	23	49	71	42	2	0	187	3.9
E	0	0	0	0	0	0	0	0	0.0
受診者数 (人)	20	873	1,937	1,402	566	37	0	4,835	100.0
有所見者数 (人)	1	252	746	654	279	19	0	1,951	40.4
有所見者率 (%)	5.0	28.9	38.5	46.6	49.3	51.4	0.0	40.4	

＜上部消化管X線検査疾患別一覧＞

＜男性＞

	～29	30～39	40～49	50～59	60～69	70～79	80～	総計
著変なし	17	1,222	1,922	1,122	668	53	0	5,004
慢性胃炎	4	207	497	503	479	36	0	1,726
びらん性胃炎	0	18	36	33	21	2	0	110
胃ポリープ	0	86	193	181	88	6	0	554
胃隆起性病変	0	7	19	18	22	3	0	69
胃潰瘍	0	0	2	5	1	0	0	8
胃潰瘍癒痕	0	12	20	26	31	5	0	94
胃粘膜下腫瘍	0	5	11	13	15	2	0	46
陰影斑(バリウム斑)	0	1	6	4	9	0	0	20
胃陥凹性病変	0	1	5	9	7	1	0	23
ひだ中断(粘膜中断)	0	1	0	1	1	0	0	3
粘膜不整	0	3	26	47	41	4	0	121
十二指腸潰瘍	0	0	0	0	0	0	0	0
十二指腸潰瘍癒痕	0	16	39	25	25	2	0	107
食道隆起性病変	0	1	4	5	5	3	0	18
総計	22	1,560	2,721	1,926	1,353	109	0	7,691

＜女性＞

	～29	30～39	40～49	50～59	60～69	70～79	80～	総計
著変なし	19	621	1,191	748	287	18	0	2,884
慢性胃炎	0	92	272	296	169	15	0	844
びらん性胃炎	0	5	20	21	12	0	0	58
胃ポリープ	1	137	413	289	79	2	0	921
胃隆起性病変	0	8	14	25	8	0	0	55
胃潰瘍	0	2	2	1	0	1	0	6
胃潰瘍癒痕	0	4	7	4	8	0	0	23

胃粘膜下腫瘍	0	5	12	20	4	2	0	43
陰影斑(バリウム斑)	0	1	2	4	3	0	0	10
胃陥凹性病変	0	2	4	3	3	0	0	12
ひだ中断(粘膜中断)	0	2	1	1	1	0	0	5
粘膜不整	0	6	19	25	13	1	0	64
十二指腸潰瘍	0	0	0	0	0	0	0	0
十二指腸潰瘍瘢痕	0	1	11	18	7	0	0	37
食道隆起性病変	0	0	1	1	3	0	0	5
総計	20	873	1,937	1,402	566	37	0	4,835

④ 施設外健診（協会けんぽ含む）

<上部消化管X線検査判定結果>

<男性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	15	509	940	807	742	343	2	3,358	53.2
B1	1	182	500	593	873	530	1	2,680	42.4
B2	0	0	0	0	0	0	0	0	0.0
C	0	0	0	0	0	0	0	0	0.0
D	0	4	25	48	135	65	0	277	4.4
E	0	0	0	0	0	0	0	0	0.0
受診者数 (人)	16	695	1,465	1,448	1,750	938	3	6,315	100.0
有所見者数 (人)	1	186	525	641	1,008	595	1	2,957	46.8
有所見者率 (%)	6.3	26.8	35.8	44.3	57.6	63.4	33.3	46.8	

<女性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	0	217	370	293	434	198	0	1,512	47.9
B1	0	68	233	324	599	324	0	1,548	49.0
B2	0	0	1	0	0	0	0	1	0.0
C	0	0	0	0	0	0	0	0	0.0
D	0	1	16	19	30	29	0	95	3.0
E	0	0	0	0	0	0	0	0	0.0
受診者数 (人)	0	286	620	636	1,063	551	0	3,156	100.0
有所見者数 (人)	0	69	250	343	629	353	0	1,644	52.1
有所見者率 (%)	0.0	24.1	40.3	53.9	59.2	64.1	0.0	52.1	

<上部消化管X線検査疾患別一覧>

<男性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	総計
著変なし	15	509	940	807	742	343	2	3,358
慢性胃炎	0	127	354	447	786	487	1	2,202
びらん性胃炎	0	14	41	72	66	37	0	230
胃ポリープ	1	42	119	116	106	50	0	434
胃隆起性病変	0	1	7	16	30	11	0	65
胃潰瘍	0	0	1	3	3	0	0	7
胃潰瘍癒痕	0	1	12	17	37	21	0	88
胃粘膜下腫瘍	0	0	2	7	16	9	0	34
陰影斑(バリウム斑)	0	1	11	9	31	15	0	67
胃陥凹性病変	0	1	0	4	6	5	0	16
ひだ中断(粘膜中断)	0	0	0	0	1	0	0	1
粘膜不整	0	0	4	6	22	13	0	45
十二指腸潰瘍	0	0	0	0	0	0	0	0
十二指腸潰瘍癒痕	0	5	20	22	33	10	0	90
食道隆起性病変	0	0	1	3	1	2	0	7
総計	16	695	1,465	1,448	1,750	938	3	6,315

<女性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	総計
著変なし	0	217	370	293	434	198	0	1,512
慢性胃炎	0	23	99	172	412	255	0	961
びらん性胃炎	0	3	20	23	54	33	0	133
胃ポリープ	0	40	123	131	147	61	0	502
胃隆起性病変	0	2	6	5	13	11	0	37
胃潰瘍	0	0	0	0	0	2	0	2
胃潰瘍癒痕	0	0	4	5	11	5	0	25
胃粘膜下腫瘍	0	0	1	3	12	3	0	19

陰影斑(バリウム斑)	0	0	5	8	3	4	0	20
胃陥凹性病変	0	1	0	2	0	0	0	3
ひだ中断(粘膜中断)	0	0	0	0	0	1	0	1
粘膜不整	0	1	0	3	4	3	0	11
十二指腸潰瘍	0	0	0	0	0	0	0	0
十二指腸潰瘍癒痕	0	1	1	9	8	7	0	26
食道隆起性病変	0	0	1	0	1	2	0	4
総計	0	286	620	636	1,063	551	0	3,156

⑤ 全国健康保険協会管掌健康保険生活習慣病予防健診[再掲]

<上部消化管X線検査判定結果>

<男性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	0	1,034	1,692	994	578	32	0	4,330	63.8
B1	0	131	262	287	169	11	0	860	12.7
B2	0	169	391	362	354	18	0	1,294	19.1
C	0	0	0	0	0	0	0	0	0.0
D	0	21	80	106	91	9	0	307	4.5
E	0	0	0	0	0	0	0	0	0.0
受診者数 (人)	0	1,355	2,425	1,749	1,192	70	0	6,791	100.0
有所見者数 (人)	0	321	733	755	614	38	0	2,461	36.2
有所見者率 (%)	0.0	23.7	30.2	43.2	51.5	54.3	0.0	36.2	

<女性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	合計 (人)	比率 (%)
A	0	446	826	502	215	7	0	1,996	58.9
B1	0	88	274	190	71	1	0	624	18.4
B2	0	81	247	218	94	2	0	642	18.9
C	0	0	0	0	0	0	0	0	0.0
D	0	14	41	47	24	0	0	126	3.7
E	0	0	0	0	0	0	0	0	0.0
受診者数 (人)	0	629	1,388	957	404	10	0	3,388	100.0
有所見者数 (人)	0	183	562	455	189	3	0	1,392	41.1
有所見者率 (%)	0.0	29.1	40.5	47.5	46.8	30.0	0.0	41.1	

<上部消化管X線検査疾患別一覧>

<男性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	総計
著変なし	0	1,034	1,692	994	578	32	0	4,330
慢性胃炎	0	204	454	477	431	24	0	1,590
びらん性胃炎	0	18	36	35	16	1	0	106
胃ポリープ	0	75	174	168	81	4	0	502
胃隆起性病変	0	6	21	14	21	2	0	64
胃潰瘍	0	0	1	5	0	0	0	6
胃潰瘍癒痕	0	10	19	24	25	3	0	81
胃粘膜下腫瘍	0	4	11	12	13	2	0	42
陰影斑(バリウム斑)	0	1	9	4	8	0	0	22
胃陥凹性病変	0	2	4	8	7	1	0	22
ひだ中断(粘膜中断)	0	0	0	1	1	0	0	2
粘膜不整	0	2	22	44	36	1	0	105
十二指腸潰瘍	0	0	0	0	0	0	0	0
十二指腸潰瘍癒痕	0	16	43	21	22	1	0	103
食道隆起性病変	0	1	2	3	4	2	0	12
総計	0	1,355	2,425	1,749	1,192	70	0	6,791

<女性>

	～29	30～39	40～49	50～59	60～69	70～79	80～	総計
著変なし	0	446	826	502	215	7	0	1,996
慢性胃炎	0	69	206	205	114	1	0	595
びらん性胃炎	0	2	11	17	9	0	0	39
胃ポリープ	0	100	309	204	57	1	0	671
胃隆起性病変	0	6	13	18	4	0	0	41
胃潰瘍	0	2	2	0	0	0	0	4
胃潰瘍癒痕	0	2	6	4	3	0	0	15
胃粘膜下腫瘍	0	3	11	14	3	1	0	32

陰影斑(バリウム斑)	0	1	2	4	2	0	0	9
胃陥凹性病変	0	0	4	2	0	0	0	6
ひだ中断(粘膜中断)	0	0	0	0	0	0	0	0
粘膜不整	0	4	13	13	10	0	0	40
十二指腸潰瘍	0	0	0	0	0	0	0	0
十二指腸潰瘍癒痕	0	0	8	15	3	0	0	26
食道隆起性病変	0	0	0	1	1	0	0	2
総計	0	629	1,388	957	404	10	0	3,388