

5. 呼吸器

(1) 胸部 X 線検査

① 1 日人間ドック

<胸部 X 線検査判定結果>

<男性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|-----------|-----|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 32 | 1,147 | 3,233 | 3,457 | 2,159 | 531 | 54 | 10,613 | 74.7 |
| B1 | 1 | 73 | 396 | 709 | 801 | 370 | 78 | 2,428 | 17.1 |
| B2 | 0 | 27 | 132 | 273 | 299 | 180 | 45 | 956 | 6.7 |
| C | 0 | 2 | 1 | 5 | 2 | 5 | 3 | 18 | 0.1 |
| D | 0 | 5 | 32 | 58 | 59 | 33 | 5 | 192 | 1.4 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数 (人) | 33 | 1,254 | 3,794 | 4,502 | 3,320 | 1,119 | 185 | 14,207 | 100.0 |
| 有所見者数 (人) | 1 | 107 | 561 | 1,045 | 1,161 | 588 | 131 | 3,594 | 25.3 |
| 有所見者率 (%) | 1.0 | 8.5 | 14.8 | 23.2 | 35.0 | 52.5 | 70.8 | 25.3 | |

<女性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|-----------|-----|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 31 | 986 | 2,990 | 3,036 | 1,580 | 380 | 31 | 9,034 | 76.8 |
| B1 | 2 | 52 | 329 | 658 | 665 | 270 | 54 | 2,030 | 17.3 |
| B2 | 0 | 13 | 76 | 133 | 184 | 81 | 22 | 509 | 4.3 |
| C | 0 | 1 | 3 | 4 | 4 | 3 | 0 | 15 | 0.1 |
| D | 0 | 10 | 25 | 65 | 57 | 16 | 7 | 180 | 1.5 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数 (人) | 33 | 1,062 | 3,423 | 3,896 | 2,490 | 750 | 114 | 11,768 | 100.0 |
| 有所見者数 (人) | 2 | 76 | 433 | 860 | 910 | 370 | 83 | 2,734 | 23.2 |
| 有所見者率 (%) | 6.1 | 7.2 | 12.6 | 22.1 | 36.5 | 49.3 | 72.8 | 23.2 | |

<胸部 X 線検査疾患別一覧>

<男性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 総計 |
|---------|-----|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 32 | 1,147 | 3,233 | 3,457 | 2,159 | 531 | 54 | 10,613 |
| 硬化性陰影 | 0 | 23 | 141 | 249 | 285 | 165 | 37 | 900 |
| 線状影 | 0 | 11 | 53 | 128 | 95 | 56 | 14 | 357 |
| 索状影 | 1 | 6 | 29 | 78 | 75 | 55 | 13 | 256 |
| 気腫性変化 | 0 | 15 | 77 | 159 | 154 | 69 | 13 | 487 |
| 限局性陰影 | 0 | 3 | 21 | 37 | 49 | 18 | 4 | 132 |
| 網状影 | 0 | 0 | 7 | 33 | 67 | 72 | 31 | 210 |
| 結節影 | 0 | 4 | 22 | 46 | 44 | 25 | 3 | 144 |
| 粒状影 | 0 | 1 | 9 | 10 | 16 | 13 | 0 | 49 |
| 浸潤影 | 0 | 0 | 1 | 3 | 3 | 6 | 0 | 13 |
| 縦隔・肺門増大 | 0 | 1 | 4 | 5 | 2 | 3 | 0 | 15 |
| 腫瘍影 | 0 | 0 | 1 | 1 | 4 | 0 | 0 | 6 |
| 胸水 | 0 | 1 | 0 | 0 | 0 | 2 | 0 | 3 |
| 気胸 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 2 |
| 受診者数（人） | 33 | 1,254 | 3,794 | 4,502 | 3,320 | 1,119 | 185 | 14,207 |

<女性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 総計 |
|---------|-----|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 31 | 986 | 2,990 | 3,036 | 1,580 | 380 | 31 | 9,034 |
| 硬化性陰影 | 0 | 23 | 162 | 365 | 337 | 112 | 28 | 1,027 |
| 線状影 | 0 | 10 | 27 | 78 | 71 | 30 | 4 | 220 |
| 索状影 | 1 | 6 | 20 | 51 | 72 | 36 | 8 | 193 |
| 気腫性変化 | 0 | 1 | 8 | 15 | 19 | 14 | 4 | 62 |
| 限局性陰影 | 0 | 6 | 17 | 31 | 42 | 7 | 4 | 107 |
| 網状影 | 0 | 0 | 1 | 8 | 18 | 10 | 4 | 41 |
| 結節影 | 0 | 4 | 21 | 30 | 27 | 12 | 3 | 97 |
| 粒状影 | 0 | 1 | 8 | 10 | 14 | 8 | 3 | 44 |
| 浸潤影 | 0 | 0 | 1 | 3 | 3 | 1 | 2 | 10 |
| 縦隔・肺門増大 | 0 | 0 | 2 | 3 | 1 | 1 | 0 | 7 |
| 腫瘍影 | 0 | 0 | 2 | 0 | 1 | 0 | 0 | 3 |
| 胸水 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 気胸 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 受診者数（人） | 33 | 1,062 | 3,423 | 3,896 | 2,490 | 750 | 114 | 11,768 |

② 2 日人間ドック

<胸部 X 線検査判定結果>

<男性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|-----------|------|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 6 | 179 | 621 | 711 | 616 | 304 | 51 | 2,488 | 74.6 |
| B1 | 1 | 9 | 70 | 125 | 194 | 161 | 58 | 618 | 18.5 |
| B2 | 0 | 1 | 14 | 34 | 61 | 64 | 20 | 194 | 5.8 |
| C | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| D | 0 | 3 | 1 | 3 | 16 | 8 | 2 | 33 | 1.0 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数 (人) | 7 | 192 | 706 | 873 | 887 | 537 | 131 | 3,333 | 100.0 |
| 有所見者数 (人) | 1 | 13 | 85 | 162 | 271 | 233 | 80 | 845 | 25.4 |
| 有所見者率 (%) | 14.3 | 6.8 | 12.0 | 18.6 | 30.6 | 43.4 | 61.1 | 25.4 | |

<女性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|-----------|------|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 3 | 35 | 191 | 261 | 236 | 167 | 9 | 902 | 73.7 |
| B1 | 1 | 0 | 25 | 42 | 71 | 86 | 34 | 259 | 21.2 |
| B2 | 0 | 2 | 1 | 9 | 17 | 18 | 6 | 53 | 4.3 |
| C | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| D | 0 | 0 | 1 | 0 | 6 | 3 | 0 | 10 | 0.8 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数 (人) | 4 | 37 | 218 | 312 | 330 | 274 | 49 | 1,224 | 100.0 |
| 有所見者数 (人) | 1 | 2 | 27 | 51 | 94 | 107 | 40 | 322 | 26.3 |
| 有所見者率 (%) | 25.0 | 5.4 | 12.4 | 16.3 | 28.5 | 39.1 | 81.6 | 26.3 | |

<胸部 X 線検査疾患別一覧>

<男性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 総計 |
|----------|-----|-------|-------|-------|-------|-------|-----|-------|
| 著変なし | 6 | 179 | 621 | 711 | 616 | 304 | 51 | 2,488 |
| 硬化性陰影 | 0 | 2 | 14 | 28 | 49 | 57 | 30 | 180 |
| 線状影 | 0 | 2 | 11 | 19 | 37 | 23 | 5 | 97 |
| 索状影 | 0 | 0 | 6 | 8 | 15 | 17 | 5 | 51 |
| 気腫性変化 | 0 | 1 | 14 | 43 | 63 | 40 | 8 | 169 |
| 限局性陰影 | 0 | 0 | 4 | 5 | 11 | 8 | 0 | 28 |
| 網状影 | 0 | 0 | 0 | 4 | 17 | 32 | 22 | 75 |
| 結節影 | 0 | 0 | 0 | 1 | 5 | 8 | 1 | 15 |
| 粒状影 | 0 | 0 | 1 | 2 | 0 | 2 | 1 | 6 |
| 浸潤影 | 0 | 0 | 1 | 0 | 1 | 1 | 0 | 3 |
| 縦隔・肺門増大 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 2 |
| 腫瘤影 | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 3 |
| 胸水 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 2 |
| 気胸 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 受診者数 (人) | 7 | 192 | 706 | 873 | 887 | 537 | 131 | 3,333 |

<女性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 総計 |
|----------|-----|-------|-------|-------|-------|-------|-----|-------|
| 著変なし | 3 | 35 | 191 | 261 | 236 | 167 | 9 | 902 |
| 硬化性陰影 | 0 | 0 | 3 | 18 | 27 | 30 | 13 | 91 |
| 線状影 | 0 | 0 | 3 | 4 | 5 | 17 | 7 | 36 |
| 索状影 | 0 | 1 | 2 | 4 | 6 | 14 | 4 | 31 |
| 気腫性変化 | 0 | 0 | 2 | 1 | 3 | 9 | 5 | 20 |
| 限局性陰影 | 0 | 0 | 1 | 1 | 4 | 2 | 0 | 8 |
| 網状影 | 0 | 1 | 0 | 0 | 4 | 5 | 2 | 12 |
| 結節影 | 0 | 0 | 1 | 2 | 5 | 1 | 0 | 9 |
| 粒状影 | 0 | 0 | 0 | 0 | 2 | 1 | 2 | 5 |
| 浸潤影 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 |
| 縦隔・肺門増大 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 腫瘤影 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| 胸水 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 気胸 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 受診者数 (人) | 4 | 37 | 218 | 312 | 330 | 274 | 49 | 1,224 |

③ 施設内健診（協会けんぽ含む）

<胸部 X 線検査判定結果>

<男性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 合計 (人) | 比率 (%) |
|----------|-------|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 2,040 | 3,307 | 3,376 | 2,298 | 1,640 | 204 | 2 | 12,867 | 85.7 |
| B1 | 40 | 139 | 319 | 317 | 389 | 77 | 0 | 1,281 | 8.5 |
| B2 | 11 | 67 | 104 | 163 | 241 | 69 | 2 | 657 | 4.4 |
| C | 0 | 1 | 0 | 0 | 2 | 0 | 0 | 3 | 0.0 |
| D | 8 | 22 | 44 | 55 | 65 | 14 | 1 | 209 | 1.4 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数(人) | 2,099 | 3,536 | 3,843 | 2,833 | 2,337 | 364 | 5 | 15,017 | 100.0 |
| 有所見者数(人) | 59 | 229 | 467 | 535 | 697 | 160 | 3 | 2,150 | 14.3 |
| 有所見者率(%) | 2.8 | 6.5 | 12.2 | 18.9 | 29.8 | 44.0 | 60.0 | 14.3 | |

<女性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 合計 (人) | 比率 (%) |
|----------|-------|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 2,436 | 2,907 | 4,042 | 2,856 | 1,145 | 129 | 7 | 13,522 | 89.2 |
| B1 | 55 | 98 | 245 | 402 | 261 | 39 | 2 | 1,102 | 7.3 |
| B2 | 6 | 24 | 72 | 126 | 98 | 24 | 6 | 356 | 2.3 |
| C | 0 | 0 | 0 | 2 | 2 | 0 | 0 | 4 | 0.0 |
| D | 5 | 20 | 39 | 64 | 37 | 7 | 0 | 172 | 1.1 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数(人) | 2,502 | 3,049 | 4,398 | 3,450 | 1,543 | 199 | 15 | 15,156 | 100.0 |
| 有所見者数(人) | 66 | 142 | 356 | 594 | 398 | 70 | 8 | 1,634 | 10.8 |
| 有所見者率(%) | 2.6 | 4.7 | 8.1 | 17.2 | 25.8 | 35.2 | 53.3 | 10.8 | |

<胸部 X 線検査疾患別一覧>

<男性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-------|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 2,040 | 3,307 | 3,376 | 2,298 | 1,640 | 204 | 2 | 12,867 |
| 硬化性陰影 | 14 | 47 | 125 | 143 | 166 | 45 | 1 | 541 |
| 線状影 | 2 | 18 | 51 | 55 | 71 | 12 | 0 | 209 |
| 索状影 | 2 | 19 | 34 | 44 | 61 | 12 | 1 | 173 |
| 気腫性変化 | 6 | 37 | 85 | 94 | 113 | 27 | 1 | 363 |
| 限局性陰影 | 1 | 4 | 18 | 15 | 24 | 6 | 0 | 68 |
| 網状影 | 0 | 7 | 8 | 13 | 62 | 27 | 3 | 120 |
| 結節影 | 2 | 10 | 20 | 24 | 35 | 3 | 1 | 95 |
| 粒状影 | 0 | 4 | 9 | 9 | 11 | 4 | 0 | 37 |
| 浸潤影 | 1 | 1 | 2 | 4 | 5 | 0 | 0 | 13 |
| 縦隔・肺門増大 | 0 | 2 | 3 | 3 | 4 | 0 | 0 | 12 |
| 腫瘍影 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 2 |
| 胸水 | 0 | 0 | 1 | 1 | 2 | | 0 | 4 |
| 気胸 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| 受診者数（人） | 2,099 | 3,536 | 3,843 | 2,833 | 2,337 | 364 | 5 | 15,017 |

<女性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-------|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 2,436 | 2,907 | 4,042 | 2,856 | 1,145 | 129 | 7 | 13,522 |
| 硬化性陰影 | 8 | 43 | 127 | 251 | 152 | 27 | 3 | 611 |
| 線状影 | 4 | 11 | 35 | 46 | 27 | 6 | 0 | 129 |
| 索状影 | 1 | 5 | 20 | 42 | 42 | 7 | 0 | 117 |
| 気腫性変化 | 1 | 2 | 9 | 12 | 14 | 3 | 0 | 41 |
| 限局性陰影 | 0 | 2 | 18 | 29 | 16 | 2 | 0 | 67 |
| 網状影 | 0 | 0 | 0 | 4 | 8 | 2 | 1 | 15 |
| 結節影 | 1 | 10 | 14 | 26 | 10 | 3 | 0 | 64 |
| 粒状影 | 0 | 3 | 8 | 21 | 17 | 2 | 1 | 52 |
| 浸潤影 | 3 | 5 | 3 | 7 | 2 | 0 | 0 | 20 |
| 縦隔・肺門増大 | 1 | 0 | 2 | 5 | 1 | 0 | 0 | 9 |
| 腫瘍影 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 2 |
| 胸水 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 気胸 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 2 |
| 受診者数（人） | 2,502 | 3,049 | 4,398 | 3,450 | 1,543 | 199 | 15 | 15,156 |

④ 施設外健診（協会けんぽ含む）

<胸部 X 線検査判定結果>

<男性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|----------|-------|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 3,769 | 4,263 | 3,533 | 3,171 | 2,566 | 1,225 | 403 | 18,930 | 86.3 |
| B1 | 46 | 145 | 269 | 412 | 486 | 353 | 223 | 1,934 | 8.8 |
| B2 | 25 | 26 | 80 | 135 | 215 | 194 | 132 | 807 | 3.7 |
| C | 0 | 2 | 0 | 1 | 2 | 1 | 1 | 7 | 0.0 |
| D | 12 | 19 | 19 | 49 | 68 | 54 | 42 | 263 | 1.2 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数(人) | 3,852 | 4,455 | 3,901 | 3,768 | 3,337 | 1,827 | 801 | 21,941 | 100.0 |
| 有所見者数(人) | 83 | 192 | 368 | 597 | 771 | 602 | 398 | 3,011 | 13.7 |
| 有所見者率(%) | 2.2 | 4.3 | 9.4 | 15.8 | 23.1 | 33.0 | 49.7 | 13.7 | |

<女性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|----------|-------|-------|-------|-------|-------|-------|------|-----------|-----------|
| A | 2,042 | 1,518 | 1,849 | 1,779 | 2,052 | 1,508 | 518 | 11,266 | 84.1 |
| B1 | 26 | 68 | 103 | 200 | 400 | 454 | 261 | 1,482 | 11.1 |
| B2 | 3 | 5 | 28 | 55 | 122 | 136 | 110 | 459 | 3.4 |
| C | 0 | 0 | 0 | 1 | 1 | 0 | 1 | 3 | 0.0 |
| D | 1 | 5 | 18 | 29 | 47 | 51 | 32 | 183 | 1.4 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数(人) | 2,072 | 1,566 | 1,998 | 2,064 | 2,622 | 2,149 | 922 | 13,393 | 100.0 |
| 有所見者数(人) | 30 | 48 | 149 | 285 | 570 | 641 | 404 | 2,127 | 15.9 |
| 有所見者率(%) | 1.4 | 3.1 | 7.5 | 13.8 | 21.7 | 29.8 | 43.8 | 15.9 | |

<胸部 X 線検査疾患別一覧>

<男性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-------|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 3,769 | 4,263 | 3,533 | 3,171 | 2,566 | 1,225 | 403 | 18,930 |
| 硬化性陰影 | 13 | 46 | 111 | 175 | 202 | 157 | 108 | 812 |
| 線状影 | 5 | 16 | 44 | 71 | 95 | 59 | 41 | 331 |
| 索状影 | 6 | 11 | 36 | 39 | 79 | 48 | 34 | 253 |
| 気腫性変化 | 11 | 17 | 56 | 128 | 108 | 52 | 30 | 402 |
| 限局性陰影 | 4 | 6 | 4 | 17 | 25 | 17 | 13 | 86 |
| 網状影 | 1 | 3 | 4 | 23 | 69 | 85 | 80 | 265 |
| 結節影 | 4 | 10 | 11 | 26 | 25 | 16 | 9 | 101 |
| 粒状影 | 0 | 3 | 2 | 11 | 9 | 7 | 3 | 35 |
| 浸潤影 | 1 | 1 | 5 | 3 | 7 | 8 | 1 | 26 |
| 縦隔・肺門増大 | 2 | 3 | 0 | 1 | 2 | 5 | 3 | 16 |
| 腫瘤影 | 0 | 1 | 1 | 0 | 2 | 2 | 2 | 8 |
| 胸水 | 0 | 2 | 0 | 0 | 5 | 2 | 6 | 15 |
| 気胸 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 受診者数（人） | 3,852 | 4,455 | 3,901 | 3,768 | 3,337 | 1,827 | 801 | 21,941 |

<女性>

| | ～29 | 30～39 | 40～49 | 50～59 | 60～69 | 70～79 | 80～ | 総計 |
|---------|-------|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 2,042 | 1,518 | 1,849 | 1,779 | 2,052 | 1,508 | 518 | 11,266 |
| 硬化性陰影 | 3 | 16 | 65 | 113 | 198 | 195 | 103 | 693 |
| 線状影 | 1 | 3 | 18 | 34 | 77 | 93 | 33 | 259 |
| 索状影 | 1 | 1 | 8 | 31 | 61 | 62 | 44 | 208 |
| 気腫性変化 | 0 | 0 | 7 | 5 | 10 | 14 | 5 | 41 |
| 限局性陰影 | 0 | 1 | 1 | 10 | 19 | 23 | 13 | 67 |
| 網状影 | 0 | 1 | 2 | 7 | 23 | 20 | 33 | 86 |
| 結節影 | 2 | 3 | 6 | 13 | 19 | 14 | 12 | 69 |
| 粒状影 | 0 | 0 | 5 | 8 | 15 | 13 | 7 | 48 |
| 浸潤影 | 0 | 1 | 5 | 2 | 7 | 6 | 3 | 24 |
| 縦隔・肺門増大 | 0 | 0 | 0 | 0 | 6 | 3 | 2 | 11 |
| 腫瘤影 | 0 | 0 | 1 | 0 | 1 | 1 | 2 | 5 |
| 胸水 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 気胸 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 |
| 受診者数（人） | 2,072 | 1,566 | 1,998 | 2,064 | 2,622 | 2,149 | 922 | 13,393 |

⑤ 全国健康保険協会管掌健康保険生活習慣病予防健診 [再掲]

<胸部 X 線検査判定結果>

<男性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|----------|-----|-------|-------|-------|-------|-------|-----|-----------|-----------|
| A | 0 | 1,700 | 3,129 | 2,169 | 1,512 | 152 | 0 | 8,662 | 82.6 |
| B1 | 0 | 74 | 295 | 317 | 362 | 56 | 0 | 1,104 | 10.5 |
| B2 | 0 | 29 | 103 | 140 | 223 | 46 | 0 | 541 | 5.2 |
| C | 0 | 0 | 0 | 0 | 3 | 0 | 0 | 3 | 0.0 |
| D | 0 | 14 | 39 | 51 | 61 | 10 | 0 | 175 | 1.7 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数(人) | 0 | 1,817 | 3,566 | 2,677 | 2,161 | 264 | 0 | 10,485 | 100.0 |
| 有所見者数(人) | 0 | 117 | 437 | 508 | 649 | 112 | 0 | 1,823 | 17.4 |
| 有所見者率(%) | 0.0 | 6.4 | 12.3 | 19.0 | 30.0 | 42.4 | 0.0 | 17.4 | |

<女性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 合計 (人) | 比率 (%) |
|----------|-----|-------|-------|-------|-------|-------|-----|-----------|-----------|
| A | 0 | 965 | 2,398 | 1,688 | 789 | 68 | 0 | 5,908 | 86.8 |
| B1 | 0 | 30 | 141 | 232 | 180 | 19 | 0 | 602 | 8.8 |
| B2 | 0 | 9 | 44 | 74 | 64 | 10 | 0 | 201 | 3.0 |
| C | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 2 | 0.0 |
| D | 0 | 5 | 23 | 39 | 25 | 2 | 0 | 94 | 1.4 |
| E | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.0 |
| 受診者数(人) | 0 | 1,009 | 2,606 | 2,035 | 1,058 | 99 | 0 | 6,807 | 100.0 |
| 有所見者数(人) | 0 | 44 | 208 | 347 | 269 | 31 | 0 | 899 | 13.2 |
| 有所見者率(%) | 0.0 | 4.4 | 8.0 | 17.1 | 25.4 | 31.3 | 0.0 | 13.2 | |

<胸部 X 線検査疾患別一覧>

<男性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 総計 |
|----------|-----|-------|-------|-------|-------|-------|-----|--------|
| 著変なし | 0 | 1,700 | 3,129 | 2,169 | 1,512 | 152 | 0 | 8,662 |
| 硬化性陰影 | 0 | 16 | 120 | 125 | 153 | 31 | 0 | 445 |
| 線状影 | 0 | 8 | 49 | 60 | 73 | 10 | 0 | 200 |
| 索状影 | 0 | 12 | 36 | 39 | 57 | 9 | 0 | 153 |
| 気腫性変化 | 0 | 18 | 83 | 92 | 110 | 18 | 0 | 321 |
| 限局性陰影 | 0 | 2 | 15 | 17 | 22 | 5 | 0 | 61 |
| 網状影 | 0 | 3 | 7 | 19 | 60 | 18 | 0 | 107 |
| 結節影 | 0 | 5 | 18 | 26 | 31 | 2 | 0 | 82 |
| 粒状影 | 0 | 4 | 5 | 5 | 9 | 4 | 0 | 27 |
| 浸潤影 | 0 | 2 | 2 | 2 | 5 | 0 | 0 | 11 |
| 縦隔・肺門増大 | 0 | 1 | 3 | 2 | 3 | 0 | 0 | 9 |
| 腫瘤影 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 胸水 | 0 | 0 | 0 | 1 | 2 | 0 | 0 | 3 |
| 気胸 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 受診者数 (人) | 0 | 1,817 | 3,566 | 2,677 | 2,161 | 264 | 0 | 10,485 |

<女性>

| | ~29 | 30~39 | 40~49 | 50~59 | 60~69 | 70~79 | 80~ | 総計 |
|----------|-----|-------|-------|-------|-------|-------|-----|-------|
| 著変なし | 0 | 965 | 2,398 | 1,688 | 789 | 68 | 0 | 5,908 |
| 硬化性陰影 | 0 | 15 | 75 | 146 | 108 | 13 | 0 | 357 |
| 線状影 | 0 | 4 | 25 | 31 | 16 | 3 | 0 | 79 |
| 索状影 | 0 | 0 | 14 | 25 | 27 | 3 | 0 | 69 |
| 気腫性変化 | 0 | 0 | 6 | 7 | 9 | 0 | 0 | 22 |
| 限局性陰影 | 0 | 0 | 9 | 16 | 11 | 1 | 0 | 37 |
| 網状影 | 0 | 0 | 0 | 2 | 7 | 1 | 0 | 10 |
| 結節影 | 0 | 4 | 7 | 14 | 6 | 0 | 0 | 31 |
| 粒状影 | 0 | 0 | 2 | 15 | 12 | 0 | 0 | 29 |
| 浸潤影 | 0 | 1 | 2 | 6 | 2 | 0 | 0 | 11 |
| 縦隔・肺門増大 | 0 | 0 | 2 | 2 | 1 | 0 | 0 | 5 |
| 腫瘤影 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胸水 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 気胸 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 受診者数 (人) | 0 | 1,009 | 2,606 | 2,035 | 1,058 | 99 | 0 | 6,807 |